

NAMI-971

THE EFFECT OF DRUGS IN ALTERING SUSCEPTIBILITY
TO MOTION SICKNESS IN AEROBATICS AND THE
SLOW ROTATION ROOM

Frederick R. Deane, Charles D. Wood, Ashton Graybiel, and Arthur C. Cawrse



JOINT REPORT



N67-26270

FACILITY FORM 602

(ACCESSION NUMBER)
14
(PAGES)
OR-84019
(NASA OR DTIC OR AD NUMBER)

(THRU)
1
(CODE)
04
(CATEGORY)

NAVAL AEROSPACE MEDICAL INSTITUTE
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

June 1966

Distribution of this document is unlimited.

Distribution of this document is unlimited.

THE EFFECT OF DRUGS IN ALTERING SUSCEPTIBILITY
TO MOTION SICKNESS IN AEROBATICS AND THE
SLOW ROTATION ROOM*

Frederick R. Deane, Charles D. Wood, Ashton Graybiel, and Arthur C. Cawrse

Bureau of Medicine and Surgery
MR005.13-6001.1.132

NASA Order R-93

Released by

Captain H. C. Hunley, MC, USN
Commanding Officer

*This research was conducted under the sponsorship of the Office of Advanced Research and Technology, National Aeronautics and Space Administration.

3 June 1966

NAVAL AEROSPACE MEDICAL INSTITUTE
NAVAL AEROSPACE MEDICAL CENTER
PENSACOLA, FLORIDA

SUMMARY PAGE

THE PROBLEM

To compare susceptibility to motion sickness and the efficacy of exemplary drugs in the prevention of motion sickness in the SRR and airsickness in aerobatics.

FINDINGS

Individual difference in drug effectiveness was demonstrated to be significant at the .01 level or better and was similar under the two conditions. Susceptibility to motion sickness in the SRR was generally a good predictor of susceptibility in aerobatics in eight subjects, but in the remaining two it was grossly in error. A combination of scopolamine and d-amphetamine was by far the most effective of the drugs tested under both conditions.

ACKNOWLEDGMENTS

The authors wish to express appreciation to LCDR R. J. Wherry, Jr., for his most valuable assistance.

INTRODUCTION

In previous reports (4,6,7) from this laboratory a standardized laboratory test procedure was described which was useful in demonstrating individual differences in susceptibility to canal sickness, a type of motion sickness experienced in a rotating environment. The relative value of exemplary drugs of different categories in reducing susceptibility to canal sickness has also been demonstrated (10, 11) by this procedure.

The basic purpose of the present experiment was to extend these studies by demonstrating whether and to what extent the findings in the Slow Rotation Room (SRR) were transferable to another force environment. This was done by comparing in the same group of subjects their susceptibility to motion sickness and the efficacy of exemplary drugs in the prevention of canal sickness in the SRR and airsickness in aerobatics.

PROCEDURE

Eighteen healthy Navy enlisted men, 18-21 years of age, assigned to the Naval Aerospace Medical Institute, were screened in a single (nonmedicated) trial of the aerobatics described below. To allow for possible habituation with continued exposure and to permit maximum separation of end points under the expected drug benefit, the ten most airsickness susceptible were chosen as study subjects.

Seven exemplary drugs were pulverized and each was placed in identical gelatin capsules, as were three individual glucose placebos (Table I).

Each "drug" was presented once to each subject in each environment according to a double blind, order-confounding, Latin Square schedule (Table II). This design was developed by R. J. Wherry, Jr., (9) for use where learning, adaptation, or residual treatment effects may be suspected to be present. Usable in squares of 4, 6, 10, 12, 16, 18, et cetera units per side, the design provides for a given drug to be both preceded and followed only once by any other drug in the total number of trials. The coding of drugs was changed between the air and SRR trials to avoid repetition of the presentation order.

The rotating force environment was generated in the Pensacola Slow Rotation Room, an essentially circular, windowless room, 15 feet in diameter and centered about the rotational axis of a human centrifuge (4). The Dial Test (6) provided a standardized series of tape-recorder paced head movements in sequences of five motions at six-second intervals. The strength of the canal sickness producing stimulus of gyroscopic forces on the vestibular system with head motion outside the plane of room rotation varies with the speed of rotation. This allows "customizing" the stimulus to subject susceptibility level.

Subject calibration at 7.5, 10, 15, and 20 rpm prior to the SRR portion of the study determined an optimal rotation speed for a given subject so that canal sickness could be expected for the nonmedicated subject in from five to fifteen sequences. When

Table I

Drugs and Dosages Used in Aerobatics and SRR Dial Test Studies

Drug	Dosage (mg)
Scopolamine with d-Amphetamine (Dexedrine)	0.6 10.0
D-Amphetamine (Dexedrine)	10.0
Cinnarazine (Mitronal)	50.0
Cyclizine (Marazine)	50.0
Meclizine (Bonamine)	50.0
Phenoxybenzamine (Dibenzylene)	20.0
Prochlorperazine (Compazine)	10.0
Placebos 1, 2, and 3 (glucose)	----

Table II

Latin Square Used To Schedule the Order of Drug Presentation in the Air or
on the SRR *

Subject	Trial									
	1	2	3	4	5	6	7	8	9	10
Hi	1	2	3	4	5	6	7	8	9	10
Le	2	4	6	8	10	1	3	5	7	9
We	3	6	9	1	4	7	10	2	5	8
Br	4	8	1	5	9	2	6	10	3	7
Du	5	10	4	9	3	8	2	7	1	6
Po	6	1	7	2	8	3	9	4	10	5
Ol	7	3	10	6	2	9	5	1	8	4
Se	8	5	2	10	7	4	1	9	6	3
Mi	9	7	5	3	1	10	8	6	4	2
Ho	10	9	8	7	6	5	4	3	2	1

*Numbers inside the square represent drugs.

it was found that no single speed could be used for all ten subjects, 10 and 20 rpm, respectively, were selected as test speeds for each of two groups of five subjects.

The air environment consisted of an observer-monitored subject seated facing aft in the rear compartment of a single-engine A1E "Skyraider" aircraft during a maximum of twenty-four aerobatic maneuvers. After a level climb to 11,000 feet in the working area under VFR conditions, the same pilot throughout performed the following sequence of maneuvers: 1-4) 3.5 G 360° steep turns alternately left and right during which the subject made 45° head tilts forward, back, and to the sides at two-second intervals; 5) nine wing-rocking and porpoising motions with the subject's gaze focused inside the aircraft; 6 and 7) wing-overs left and right; 8 and 9) aileron rolls left and right; 10 and 11) barrel rolls left and right; 12) a split-S with 4 G pullout. During the final six maneuvers, the subject's attention was focused on reading material held in his lap. Maneuvers 1-4 required about 6,000 feet, 5-11 were at a nearly constant altitude, and 12 required 3,000 feet. The entire sequence lasted fifteen minutes and was repeated once after return to 11,000 feet if an end point had not been reached. The same stimulus strength in the air was experienced by all ten subjects.

The end point for diagnosis of motion sickness, based on the criteria shown in Table III was Malaise III (11). When this stage was reached, straight and level flight was immediately resumed or, in the SRR, the head was fixed and rotation stopped. A maximum of twenty-four maneuvers in the air or sixty sequences (300 head movements) on the SRR was allowed if Malaise III was not reached. Subjects were instructed to report the same level of discomfort and a similar end point in each trial. Scoring was in terms of the number of maneuvers (twenty-four maximum) or sequences (sixty maximum) completed in reaching Malaise III.

Data were analyzed in two ways. A rank order of drug effectiveness (the average of the scores of ten subjects for a given drug) and of subject susceptibility (the average of the three placebo scores for a given subject) was obtained for each environment. During the SRR nonmedicated calibration runs, each of the five men later run at 20 rpm in the drug trials was considerably less susceptible than all five men later run at 10 rpm. In assigning a rank order of subject susceptibility, it was assumed that this was also true in the placebo trials. A rank-difference coefficient of correlation for subject susceptibility and drug effectiveness in the two environments was obtained for significance testing.

In addition, since the aerobatic maneuvers were assumed to be comparable, the data were computer analyzed for each environment separately by the analysis of variance. A graphical comparison of drug effectiveness was made, expressing it as the percentage increase in number of maneuvers or sequences for a given drug over that with Placebo 1.

Table III

IMPORTANT VESTIBULAR SYMPTOMS* USED IN DIAGNOSTIC CATEGORIZATION			
PATHOG- NOMONIC	MAJOR	MINOR	DIAGNOSTIC TERMS
VOMIT- ING	RETCHING		VESTIBULAR SICKNESS:
	NAUSEA III OR II	NAUSEA I	VOMITING OR TWO MAJOR SYM. OR ONE MAJOR & TWO MINOR
	INC. SALIV. III OR II	INC. SALIV. I	MAL AISE III : #
	PALLOR III	PALLOR II	ONE MAJOR OR TWO MINOR OR ONE MINOR & TWO OTHER
	COLD SWEAT III	COLD SWEAT II	MAL AISE I :
	DROWSINESS III	DROWSINESS II	ANY SUBJ. SYM. OR ANY SIGN USUALLY ASSOC. WITH SUBJ. SYM.
			MAL AISE II : ALL OTHER

* IN RARE INSTANCES OTHER SYMPTOMS QUALIFY

MAL AISE III WAS USED AS AN END POINT IN THE PRESENT STUDY

RESULTS

Rank order of subject susceptibility in each environment is shown in Table IV. While those subjects more susceptible to sickness in the air were generally so on the SRR, exceptions occurred, e.g., subjects Le and Po. The rank-difference coefficient of correlation was not significant at the .05 level.

Rank order of drug effectiveness in each environment is shown in Table V. The rank-difference coefficient of correlation for the two conditions was significant at the .05 level.

In the analysis of variance, drug effect was significant at the .01 and .001 levels in the air and SRR, respectively. Subject susceptibility in the air was significant at the .001 level. However, while a given subject was rotated at a constant speed for the drug trials, allowing a rank order of susceptibility to be assigned based on the calibration run data, subject susceptibility on the SRR was not meaningful since different subjects were rotated at two different speeds for the drug trials.

Trial order was not significant in either condition; habituation was not shown under these exposure conditions.

Graphical comparison of drug effectiveness in the air and in the SRR is shown in Figure 1. Those drugs most effective in the air were generally most effective on the SRR. The three separately treated placebos were very closely grouped (as to effectiveness) in both environments, indicating their validity for control purposes.

The scopolamine/d-amphetamine mixture was by far the most effective in both conditions. Compared to Placebo 1, the mixture increased tolerance in the air by 78 per cent and in the SRR by 188 per cent. No drug was more effective than scopolamine with d-amphetamine in seven subjects in the air and in eight in the SRR.

D-amphetamine alone was the second most effective drug in both environments, being the most effective (alone or one of two) for three of the ten subjects in the air and one on the SRR. In the air, cyclizine, cinnarazine, and meclizine provided the best protection for one subject each; on the SRR, cinnarazine was the most effective in two subjects. Dibenzylene had an unfavorable effect in both environments while meclizine apparently led to an increase in susceptibility in the SRR.

Side effects were minor and randomly dispersed with respect to drugs and subjects; scopolamine plus d-amphetamine did not produce symptoms significantly different or greater than placebos.

DISCUSSION

In the present study it was not possible to equate completely one aerobic maneuver with all others, although they were roughly similar in stress duration, type,

Table IV

Rank Order of Subject Susceptibility in the Air and SRR*

	Ho	Le	Mi	Br	Subject Ol	Se	Po	We	Du	Hi
Air	1.5	1.5	3.0	4.0	5.5	5.5	7.5	7.5	9.0	10.0
SRR	2.0	9.0	3.0	6.0	5.0	4.0	1.0	7.0	8.0	10.0

* 1 = Most sensitive; 10 = least sensitive

Table V

Rank Order of Drug Effectiveness in the Air and SRR*

Drug	Air	SRR
Scopolamine and d-Amphetamine	1	1
d-Amphetamine	2	2
Prochlorperazine (Compazine)	3	7
Cyclizine (Marazine)	4	4
Placebo 2	5	5
Meclizine (Bonamine)	6	10
Placebo 3	7	6
Cinnarazine (Mitronal)	8	3
Placebo 1	9	8
Phenoxybenzamine (Dibenzylene)	10	9

* 1 = Most effective; 10 = least effective

DURATION OF MANEUVERS CAUSING MALAISE III, PLACEBO I = 100%

8




Figure 1

Comparison of Antimotion Sickness Drug Effectiveness in 10 Subjects in the Air and on the SRR Dial Test

and strength. Hence, in the strictest sense, only a rank order comparison of drug effectiveness and of subject susceptibility was possible. Considering rank-difference coefficients of correlation at the .05 level, the SRR Dial Test provided valid information regarding the effectiveness of exemplary drugs under the air conditions of the experiment.

Additional data are provided by assuming that one aerobic maneuver was essentially equivalent to every other, thereby making the number of maneuvers a linear function of stress provided. With this assumption, drug effectiveness in the air and SRR was significant at the .01 and .001 levels, respectively, and may be compared as in Figure 1. The essentially linear placement of points supports the application of the SRR drug data to aerobic conditions as does the close grouping of the three placebos treated individually.

The SRR Dial Test with the Malaise III end point has many advantages. It provides a standardized, reproducible stimulus which may be widely varied in strength, ended immediately by head fixation and cessation of rotation, and repeated with a minimum of adaptation or the reduction of subject motivation due to emesis, allowing a subject to serve as his own control. It appears that this laboratory test procedure is more useful than some other devices, e.g., swings, which have been somewhat doubtful indicators of drug effectiveness under field conditions (2). Extension of this test procedure to other drugs with extrapolation to air conditions seems warranted, allowing great savings in experimental outlay while providing field applicable information.

While this study confirms that susceptibility to SRR canal sickness generally predicts sensitivity to airsickness under aerobic conditions (7), a correlation which is not significant at the .05 level was found. Other studies (2) have shown that susceptibility in one motion sickness producing environment is not an infallible predictor for another. One might speculate that variability in flight experience and anxiety levels, as well as the differences in the stimuli provided in the two environments, may be contributory.

Within the exemplary drugs studied, scopolamine, a parasympatholytic, combined with d-amphetamine, a sympathomimetic, was the most effective medication by a wide margin; d-amphetamine alone was second in both environments, confirming studies in this laboratory (10,11) and others (1,2,5,8,12). Dibenzylene, a sympatholytic, was shown to increase motion sickness susceptibility slightly as reported by Chinn et al.(3). These findings harken to earlier theories (5) which noted that motion sickness symptoms resemble parasympathetic overactivity, may be reproduced by anticholinesterases such as physostigmine, and might be prevented or reduced by drugs shifting the balance toward the sympathetic. It is difficult to reconcile a single centrally active mechanism with these facts since scopolamine has a depressant action and both d-amphetamine and Dibenzylene are stimulants. However, Dibenzylene has the reported central side effect of producing nausea and emesis in nonmoving environments.

Compazine, a phenothiazine tranquillizer, was moderately effective in the aerobatic portion (third, next to d-amphetamine), probably through reduction of anxiety. In the SRR it was ineffective as in previous studies (10, 11).

The three antihistamines, cyclizine, cinnarazine, and meclizine, provided contradictory results. Cyclizine was moderately effective in both experimental environments while the related cinnarazine was effective in the SRR but not in the air. Meclizine provided a contrast to previous findings (10-12) in that it was ineffective in the air and increased susceptibility in the SRR. However, subject Hi, usually the most motion sickness resistant subject and high scorer, had an atypical, extremely low score and probably during his SRR meclizine trial, a mild gastroenteritis was developing. Recalculation of rank order of drug effects on the SRR, eliminating Hi's scores, places meclizine in a somewhat effective range (sixth) while changing other positions little. Aside from speculation as to the quality of meclizine stock used, no other explanation can be given.

This study supports the contention that the SRR Dial Test is an effective standardized laboratory procedure for the investigation of antimotion sickness drugs and that findings, generally, may be extrapolated to air conditions. Further anti-motion sickness drug screening, the establishment of the optimal scopolamine and d-amphetamine combination, and comparison of drug effectiveness at sea and on other motion simulators are suggested.

REFERENCES

1. Blackham, R. J., Seasickness. Brit. med. J., 2:163-167, 1939.
2. Chinn, H. I., and Smith, P. K., Motion sickness. Pharmacol. Rev., 7:33-82, 1955.
3. Chinn, H. I., Handford, S. W., Smith, P. K., Cone, T. E., Jr., Redmond, R. F., Maloney, J. V., and Smyth, C. McC., Evaluation of some drugs in seasickness. J. Pharmacol., 108:69-79, 1953.
4. Graybiel, A., Clark, B., and Zarriello, J. J., Observations on human subjects living in a slow rotation room for periods of two days. Arch. Neurol., 3:55-73, 1960.
5. Hill, J., Benzedrine in seasickness. Brit. med. J., 2:1109-1112, 1937.
6. Kennedy, R. S., and Graybiel, A., The Dial Test: A standardized procedure for the experimental production of canal sickness symptomatology in a rotating environment. NSAM-930. NASA R-93. Pensacola, Fla.: Naval School of Aviation Medicine, 1965.
7. Kennedy, R. S., and Graybiel, A., Validity of tests of canal sickness in predicting susceptibility to airsickness and seasickness. Aerospace Med., 33:935-938, 1962.
8. Keevil, J. J., Benzedrine in seasickness. J. Roy. Nav. Med. Serv., 24:219-222, 1938.
9. Wherry, R. J., Jr., Personal communication.
10. Wood, C. D., Graybiel, A., and Kennedy, R. S., A comparison of effectiveness of some antimotion sickness drugs using recommended and larger than recommended doses as tested in the slow rotation room. Aerospace Med., 37:259-262, 1966.
11. Wood, C. D., Graybiel, A., and McDonough, R. G., Human centrifuge studies on the relative effectiveness of some antimotion sickness drugs. Aerospace Med., 37:187-190, 1966.
12. Wood, C. D., Kennedy, R. S., and Graybiel, A., Review of motion sickness drugs from 1954-1964. Aerospace Med., 36:1-4, 1965.

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Naval Aerospace Medical Institute Pensacola, Florida		2a. REPORT SECURITY CLASSIFICATION Unclassified	
		2b. GROUP N/A	
3. REPORT TITLE THE EFFECT OF DRUGS IN ALTERING SUSCEPTIBILITY TO MOTION SICKNESS IN AEROBATICS AND THE SLOW ROTATION ROOM			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) N/A			
5. AUTHOR(S) (Last name, first name, initial) Deane, Frederick R., LT, MC, USN, Graybiel, Ashton, CAPT, MC, USN, and Cawrse, Arthur C., LT, USN			
6. REPORT DATE 3 June 1966		7a. TOTAL NO. OF PAGES 11	7b. NO. OF REFS 12
8a. CONTRACT OR GRANT NO. NASA Order R-93		9a. ORIGINATOR'S REPORT NUMBER(S) NAMI-971	
b. PROJECT NO. MR005.13-6001.1			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.		132	
10. AVAILABILITY/LIMITATION NOTICES Distribution of this document is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
13. ABSTRACT Seven exemplary antemotion sickness drugs and three "individually treated" placebos were investigated in ten men during twenty-four aerobatic maneuvers in an A1E "Skyraider" aircraft and in performance of the Pensacola Slow Rotation Room Dial Test. The rank order of drug effectiveness and of subject susceptibility under each condition was determined and compared. Individual difference in drug effectiveness was significant at the .01 level or better and was similar under the two conditions. Susceptibility to motion sickness in the SRR was generally a good predictor of susceptibility in aerobatics in eight subjects, but in the remaining two it was grossly in error. A combination of scopolamine and d-amphetamine was by far the most effective of the drugs tested under both conditions.			

Unclassified

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Motion sickness Pharmacology Stress						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year; or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (*paying for*) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.

Unclassified

Security Classification